

AD-A117 473

ARMY INST OF DENTAL RESEARCH WASHINGTON DC  
CONTROLLED RELEASE OF ANTIBIOTICS FROM BIODEGRADABLE MICROCAPSU--ETC(U)  
JUN 82 J A SETTERSTROM, T R TICE, D H LEWIS

F/G 6/15

UNCLASSIFIED

NL

1 1/4  
A 1/2  
10 1/4




END

DATE

FILED

09:82

DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS 1963-A

18 JUN 1982

SETTERSTROM, TICE, LEWIS, and MEYERS

①

AD A117473

CONTROLLED RELEASE OF ANTIBIOTICS FROM BIODEGRADABLE  
MICROCAPSULES FOR WOUND INFECTION CONTROL

JEAN A. SETTERSTROM, Ph.D.,\* U. S. ARMY INSTITUTE OF DENTAL RESEARCH,  
WALTER REED ARMY MEDICAL CENTER, WASHINGTON, D. C. 20012  
THOMAS R. TICE, Ph.D., DANNY H. LEWIS, Ph.D., and WILLIAM E. MEYERS, Ph.D.,  
SOUTHERN RESEARCH INSTITUTE, BIRMINGHAM, ALABAMA 35255

INTRODUCTION

Improved methods to prevent and treat infection in contaminated wounds following traumatic injury are of military significance. Combat wounds are characterized by a high incidence of infection due to the inevitable presence of devitalized tissue and foreign body contaminants. Infection in these wounds is most effectively suppressed the first four hours after injury when an increased blood supply transports phagocytic cells and bactericidal factors to the injured area.(1) After four hours, vascular permeability decreases sharply and a fibrous coagulum forms at the base of the granulating wound which blocks antibiotics in the serum from penetrating the area.(2) To control infection, systemic antibiotics must be administered early when circulation is optimal. If treatment is delayed, a milieu for bacterial growth is established and complications associated with established infections occur.(3) Once infections are established, it becomes difficult to administer systemic antibiotics for an extended time, at levels that are safe, yet effective at the wound site. An ideal mode of antibiotic delivery to a contaminated wound would have the following characteristics: (1) local application in a single dose would be possible; (2) it would provide an initial burst of antibiotic for immediate tissue perfusion; and (3) it would provide a sustained and effective tissue level at the wound site. An antibiotic delivery system that fulfills these criteria is currently being developed and evaluated.

Unless topically active, drugs are distributed through the body in plasma, and the amount of drug that hits its target, is only a small part of the total drug in the body. This ineffective use of the drug is compounded in the trauma patient by hypovolemic shock, which produces a low perfusion rate of blood to tissues.(4) The ability to administer controlled quantities of a drug to a local area for a sustained time, via a biocompatible, biodegradable vehicle, provides an opportunity to evaluate a potentially advantageous method of antibiotic delivery to contaminated

DTIC FILE COPY

DTIC  
ELECTE  
S JUL 22 1982 D  
B

DISTRIBUTION STATEMENT A

Approved for public release  
Distribution Unlimited

82 07 19 21 9

page /

or infected tissue. Historically, the first success with topical antibiotics occurred at Pearl Harbor in 1941 when promptly debrided wounds, which were sprinkled with sulfanilamide, yielded low infection rates. Following inappropriate use, however, concretions of sulfa formed in the tissue causing infection rates to rise.(5) Such failures in the use of topical antibiotics are now believed to have occurred because improper drugs were used, or the method of administration failed to release drugs in an effective concentration. In recent years, topical antibiotics have been shown to be lifesaving in the prophylaxis and treatment of burn infections because the antibiotic can be applied repeatedly to the infected area, assuring a constant source of drug. The role of topical antibiotics in deep tissue wounds, however, has had less dramatic success. In contrast to burn therapy, an infected, deep tissue wound cannot be repeatedly treated with topical antibiotics unless the infected area is surgically re-exposed. The effect of single doses of antibiotics, in the form of powders, sprays, or lavages, are short-term because they are rapidly removed from the site and excreted. Nevertheless, studies have shown repeatedly that when risk of infection is high, topical antibiotics applied in this way do have an effect in reducing infection rates.(6-9) Until the recent development of sustained release systems, the opportunity to locally release drugs in tissues for sustained periods was not possible. Consequently, the theoretical advantages of such a system could not be evaluated.

It is the goal of this study to develop microcapsules that slowly release effective therapeutic doses of antibiotics in a wound over a 14-day period, by which time the microcapsules will have been biodegraded. It is the purpose of this paper to report the *in vivo* results obtained using recently formulated prototype microcapsules.

[illegible]

1  
 2  
 3  
 4  
 5  
 6  
 7  
 8  
 9  
 10  
 11  
 12  
 13  
 14  
 15  
 16  
 17  
 18  
 19  
 20  
 21  
 22  
 23  
 24  
 25  
 26  
 27  
 28  
 29  
 30  
 31  
 32  
 33  
 34  
 35  
 36  
 37  
 38  
 39  
 40  
 41  
 42  
 43  
 44  
 45  
 46  
 47  
 48  
 49  
 50  
 51  
 52  
 53  
 54  
 55  
 56  
 57  
 58  
 59  
 60  
 61  
 62  
 63  
 64  
 65  
 66  
 67  
 68  
 69  
 70  
 71  
 72  
 73  
 74  
 75  
 76  
 77  
 78  
 79  
 80  
 81  
 82  
 83  
 84  
 85  
 86  
 87  
 88  
 89  
 90  
 91  
 92  
 93  
 94  
 95  
 96  
 97  
 98  
 99  
 100  
 101  
 102  
 103  
 104  
 105  
 106  
 107  
 108  
 109  
 110  
 111  
 112  
 113  
 114  
 115  
 116  
 117  
 118  
 119  
 120  
 121  
 122  
 123  
 124  
 125  
 126  
 127  
 128  
 129  
 130  
 131  
 132  
 133  
 134  
 135  
 136  
 137  
 138  
 139  
 140  
 141  
 142  
 143  
 144  
 145  
 146  
 147  
 148  
 149  
 150  
 151  
 152  
 153  
 154  
 155  
 156  
 157  
 158  
 159  
 160  
 161  
 162  
 163  
 164  
 165  
 166  
 167  
 168  
 169  
 170  
 171  
 172  
 173  
 174  
 175  
 176  
 177  
 178  
 179  
 180  
 181  
 182  
 183  
 184  
 185  
 186  
 187  
 188  
 189  
 190  
 191  
 192  
 193  
 194  
 195  
 196  
 197  
 198  
 199  
 200  
 201  
 202  
 203  
 204  
 205  
 206  
 207  
 208  
 209  
 210  
 211  
 212  
 213  
 214  
 215  
 216  
 217  
 218  
 219  
 220  
 221  
 222  
 223  
 224  
 225  
 226  
 227  
 228  
 229  
 230  
 231  
 232  
 233  
 234  
 235  
 236  
 237  
 238  
 239  
 240  
 241  
 242  
 243  
 244  
 245  
 246  
 247  
 248  
 249  
 250  
 251  
 252  
 253  
 254  
 255  
 256  
 257  
 258  
 259  
 260  
 261  
 262  
 263  
 264  
 265  
 266  
 267  
 268  
 269  
 270  
 271  
 272  
 273  
 274  
 275  
 276  
 277  
 278  
 279  
 280  
 281  
 282  
 283  
 284  
 285  
 286  
 287  
 288  
 289  
 290  
 291  
 292  
 293  
 294  
 295  
 296  
 297  
 298  
 299  
 300  
 301  
 302  
 303  
 304  
 305  
 306  
 307  
 308  
 309  
 310  
 311  
 312  
 313  
 314  
 315  
 316  
 317  
 318  
 319  
 320  
 321  
 322  
 323  
 324  
 325  
 326  
 327  
 328  
 329  
 330  
 331  
 332  
 333  
 334  
 335  
 336  
 337  
 338  
 339  
 340  
 341  
 342  
 343  
 344  
 345  
 346  
 347  
 348  
 349  
 350  
 351  
 352  
 353  
 354  
 355  
 356  
 357  
 358  
 359  
 360  
 361  
 362  
 363  
 364  
 365  
 366  
 367  
 368  
 369  
 370  
 371  
 372  
 373  
 374  
 375  
 376  
 377  
 378  
 379  
 380  
 381  
 382  
 383  
 384  
 385  
 386  
 387  
 388  
 389  
 390  
 391  
 392  
 393  
 394  
 395  
 396  
 397  
 398  
 399  
 400  
 401  
 402  
 403  
 404  
 405  
 406  
 407  
 408  
 409  
 410  
 411  
 412  
 413  
 414  
 415  
 416  
 417  
 418  
 419  
 420  
 421  
 422  
 423  
 424  
 425  
 426  
 427  
 428  
 429  
 430  
 431  
 432  
 433  
 434  
 435  
 436  
 437  
 438  
 439  
 440  
 441  
 442  
 443  
 444  
 445  
 446  
 447  
 448  
 449  
 450  
 451  
 452  
 453  
 454  
 455  
 456  
 457  
 458  
 459  
 460  
 461  
 462  
 463  
 464  
 465  
 466  
 467  
 468  
 469  
 470  
 471  
 472  
 473  
 474  
 475  
 476  
 477  
 478  
 479  
 480  
 481  
 482  
 483  
 484  
 485  
 486  
 487  
 488  
 489  
 490  
 491  
 492  
 493  
 494  
 495  
 496  
 497  
 498  
 499  
 500  
 501  
 502  
 503  
 504  
 505  
 506  
 507  
 508  
 509  
 510  
 511  
 512  
 513  
 514  
 515  
 516  
 517  
 518  
 519  
 520  
 521  
 522  
 523  
 524  
 525

the drug vehicle. This copolymer is ideally suited for *in vivo* drug release since it elicits a minimal inflammatory response, is biologically compatible, and degrades under physiologic conditions.(10) The degradation products are nontoxic and readily metabolized. All microcapsules currently formulated exist as free-flowing microspheres (<250 microns in diameter) consisting of ampicillin anhydrate coated with a poly (DL-lactide-co-glycolide) excipient having a lactide:glycolide ratio of 68:32. Microcapsules of this size can be administered directly to a wound by a shaker-type dispenser or aerosol spray. The rate of biodegradation is controllable because it is related to the molar ratios of the constituent polymers and to the surface area of the microcapsules.

## MATERIALS AND METHODS

### Microencapsulation Process

Both solvent evaporation and phase separation methods were used in formulating the microcapsules.(11) The microencapsulation process will be described in detail in a subsequent publication, therefore, only a brief description of the process follows. In the solvent evaporation process, ampicillin anhydrate (Bristol Laboratories) was suspended in a polymer solution prepared by dissolving poly (DL-lactide-co-glycolide) in an acetone and methylene chloride solvent. This mixture was then added to a stirred aqueous solution of poly (vinyl alcohol) to form a stable oil-in-water emulsion. The oil microdroplets which formed contain drug, polymer, and polymer solvent. Removal of the polymer solvent by evaporation resulted in solid microcapsules. The poly (DL-lactide-co-glycolide) was synthesized from DL-lactide and glycolic acid. In the phase separation method the antibiotic was suspended in a dilute polymer solution. A non-solvent for both polymer and drug was added to precipitate the polymer onto suspended drug particles in order to produce the microcapsules.

### Analytical Procedures Used to Characterize the Microcapsules

Before *in vivo* testing of the microcapsules, the antibiotic content (core load) and *in vitro* release kinetics were evaluated. The core loads were determined by dissolving milligram quantities of microcapsules in methylene chloride and extracting the antibiotic with four volumes of water. The drug dissolved in the water was assayed by direct spectrophotometry, ninhydrin-based colorimetry, or microbiologic techniques. The study of the *in vitro* release rate of the antibiotic was performed by placing known amounts of microcapsules in flasks containing deionized water and agitating at 37°C. Aliquots periodically removed from the receiving fluid were assayed for drug content. A reagent prepared as a ninhydrin-hydrinatin solution was used in a colorimetric assay to evaluate the antibiotic content of the receiving fluid. Reactions of this reagent

with antibiotic solutions of various concentrations developed a color proportional in intensity to the antibiotic content.

Using both microencapsulation processes,  $^{14}\text{C}$ -labeled ampicillin anhydrate microcapsules were synthesized. (12) These radiolabeled microcapsules provided an accurate method for determining ampicillin core loadings and *in vitro* release profiles.

#### In Vivo Evaluation of Microcapsules

Ampicillin microcapsules formulated by both the solvent evaporation and phase separation processes were evaluated *in vivo* to determine the effect of the locally released drug on artificially induced wound infections. Experiments were performed on male Walter Reed strain albino rats, weighing 250-300 grams, that were anesthetized with sodium pentobarbital. The right hind leg was razor-shaved, scrubbed with Betadine, and swabbed with 70% isopropyl alcohol. A wound 2.5 to 3.0 cm in length and 5.0 mm deep was made in the thigh muscle, after which, 0.2 g of sterile dirt was added. The muscles were traumatized by pinching uniformly with tissue forceps, and inoculated with known quantities of *Staphylococcus aureus* ATCC 6538P and *Streptococcus pyogenes* ATCC 19615. The artificially contaminated wounds were treated by layering sterile, preweighed amounts of microencapsulated antibiotic directly on the wound and suturing the skin closed. Groups of animals with treated wounds (ampicillin-loaded microcapsules), untreated wounds, wounds packed with unloaded microcapsules, and wounds packed with unencapsulated antibiotic were evaluated at daily intervals.

After the effectiveness of microcapsules A681-31-1 was established, a dose-response experiment was performed wherein doses of microcapsules ranging from 0.5 to 0.05 g were applied to wounds. Sixty-eight rats were divided into five groups (A through E); four groups of 15 rats and one group of 8 rats. All rats were infected on the same day with the same quantitated bacterial suspension to assure uniform inoculum in all wounds. Wounds in the group of 8 rats (Group A) were treated with 0.5 g of ampicillin microcapsules. Rats in Groups B, C, and D were treated with 0.25, 0.10, and 0.05 g, respectively. Rats in Group E remained untreated. Bacterial counts were performed on homogenized, preweighed tissue that was removed aseptically from the wound sites. Tissue from varying distances around the wound site and serum removed by cardiac puncture were assayed for antibiotic content. This was performed by placing disks saturated with known quantities of serum or tissue homogenates on the surface of Muller-Hinton agar previously seeded with standardized amounts of *Sarcina lutea* ATCC 9341. Following incubation at 37°C, inhibition zones were measured. Freshly diluted stock solutions containing known quantities of ampicillin anhydrate served as standards. Diameters of the inhibition zones were converted to antibiotic concentrations using standard curves generated by plotting the logarithm of the drug concentration against the

Insert last  
name of au-  
thor(s) here

Start here  
for all  
pages  
after the  
first

First  
page type  
number

Author  
affiliation  
city, state

Page line

Continuing/  
page 5

SETTERSTROM, TICE, LEWIS and MEYERS

zone diameters.

## RESULTS AND DISCUSSION

The ampicillin microcapsules evaluated *in vivo* are listed in Table 1. The doses applied to each wound and the ampicillin core loadings (wt%) for each batch of microcapsules evaluated are shown in Table 2. With time, all microcapsules tested effectively reduced bacterial counts in contaminated wounds. However, microcapsules produced by the phase separation process were optimally effective in eliminating infection. An infection was considered eliminated when the wound site was bacteria free at 14 days. When 0.5 g of microcapsules (A382-140-1) were applied to wounds infected with *Staphylococcus aureus* and *Streptococcus pyogenes*, 60% of the wounds were sterile by 14 days. The remaining wounds were infected with *Staphylococcus aureus* only, since *Streptococcus pyogenes* was eliminated from all wounds by 48 hours. Wounds treated with an amount of powdered ampicillin equivalent to the core load amount, but not encapsulated within the DL-PLGA microcapsules, remained infected. Although 40% of the wounds remained contaminated at 14 days, the bacterial counts for these wounds were significantly lower than those observed for wounds treated with topical ampicillin powder or unloaded microcapsules (Table 3).

Results of the dose-response experiment performed to determine the smallest effective dose for microcapsules A681-31-1 are shown in Table 4. The bacterial counts listed in this table are for *Staphylococcus aureus* only, since all doses of microcapsules (A681-31-1) also eliminated *Streptococcus pyogenes* by 48 hours. At 7 days the wounds treated with encapsulated ampicillin remained infected with *Staphylococcus aureus*. By 14 days all wounds treated with encapsulated ampicillin were sterile; whereas, all untreated wounds remained infected. Doses of encapsulated ampicillin as small as 0.05 g per wound successfully eliminated *Staphylococcus aureus*. Based on the ampicillin core load, this quantity of microcapsule (0.05 g) contained approximately 9.05 mg (9050 µg) of ampicillin. If released uniformly over 14 days approximately 646 µg of ampicillin would be released into the wound. Kinetic studies of ampicillin released from C<sup>14</sup> labeled ampicillin anhydrate microcapsules formulated by the phase separation process showed that only 60% of the total reservoir of ampicillin was released by 14 days. Considering this, approximately 387 µg of ampicillin was available for release per day. The amounts of ampicillin detected in muscle tissue removed from wounds treated with 0.05 g of microcapsule were 54, 60, and 21 µg/g of tissue at 2, 7, and 14 days respectively. This amount is theoretically more than adequate to effectively control the growth of *Staphylococcus aureus* since the minimal inhibitory concentration sufficient to kill 95% of all strains *in vitro* is 0.5 µg/ml. An *in vitro* ampicillin level as low as 0.05 µg/ml, or 10 times less, will inhibit 97% of all strains of *Streptococcus pyogenes*.

STAMP Security  
Classification  
here

STAMP Security Classification  
here

SETTERSTROM, TICE, LEWIS, and MEYERS

Microcapsules produced by the solvent evaporation process had low ampicillin core loadings (3.0-4.5 wt%). Kinetic studies showing the *in vitro* release of ampicillin from these microcapsules indicated that only 40% of the ampicillin was released by 14 days.(13) Nevertheless, these microcapsules eliminated infections and decreased bacterial counts when applied to infected wounds. However, even though large doses were applied (1.0-0.7 g microcapsule/wound) ampicillin was not detected in serum. Rats treated with high core loaded microcapsules produced by phase separation (A681-31-1) at a dose of 0.25 g/wound, had serum ampicillin levels present the first 4 days after treatment (Figure 2). Those treated with 0.5 g per wound had serum ampicillin levels 7 days post-treatment. No serum ampicillin was detected in rats treated with 0.10 g of microcapsule per wound or less.

The microcapsules currently evaluated have been formulated with 68:32 poly(DL-lactide-co-glycolide). The bioresorption time for unsterilized microparticles of that polymer is approximately 3-4 months *in vivo*. These microcapsules deliver drug at an efficacious rate over a target period of 2 weeks; however, they also release drug at a very slow rate for over 30 days following this initial 2 week period. Once infections in wounds are eliminated, the antibiotic and microcapsules are no longer wanted in the tissue. Therefore, new microcapsules consisting of 50:50 poly(DL-lactide-co-glycolide) are being formulated. It is expected that these microcapsules will begin to biodegrade immediately following administration, and be completely degraded by 30 days. The more rapid biodegradation should reduce or eliminate the slow release of ampicillin that is occurring past 14 days.

SETTERSTROM, TICE, LEWIS, and MEYERS

In conducting the research described in this report, the investigators adhered to the "Guide for the Care and Use of Laboratory Animals" as promulgated by the Committee on the Guide for Laboratory Animal Facilities and Care, of the Institute of Laboratory Animal Resources, National Academy of Sciences, National Research Council.

Commercial materials and equipment are identified in this report to specify the investigative procedures. Such identification does not imply recommendation or endorsement or that the materials and equipment are necessarily the best available for the purpose. Furthermore, the opinions expressed herein are those of the author and are not to be construed as those of the U. S. Army Medical Department.

#### ACKNOWLEDGMENTS

The authors are grateful to Maria A. Derevjanik and Jerry S. G. Harrington for their excellent technical assistance.

THE UNIVERSITY OF CHICAGO LIBRARY

1961

TABLE 1. *IN VIVO* AMPICILLIN MICROCAPSULES EVALUATED

CAPSULE NUMBER	MICROENCAPSULATION PROCESS	MICROCAPSULE SIZE, $\mu$
9306-142-1	Solvent Evaporation	<250
A026-62-1	Solvent Evaporation	63-250
A382-140-1	Phase Separation	45-106
A681-31-1	Phase Separation	45-106

TABLE 2. AMPICILLIN CONTENT AND DOSE OF MICROCAPSULES APPLIED TO WOUNDS

<i>IN VIVO</i> EXPERIMENT	CAPSULE NUMBER	MICROCAPSULE DOSE/WOUND (AMPICILLIN EQUIVALENT)	ANTIBIOTIC CORE LOAD (WT%)
(EFFICACY)	9306-142-1	1.0 g (30.9 mg)	3
(EFFICACY)	A026-62-1	0.7 g (32.9 mg)	4.5
(EFFICACY)	A382-140-1	0.5 g (113 mg)	18.5
(DOSE-RESPONSE)	A681-31-1	0.50 g (110 mg) 0.25 g (45.25 mg) 0.10 g (18.10 mg) 0.05 g (9.05 mg)	18.1

Security Classification  
None

SETTERSTROM, TICE, LEWIS, and MEYERS

TABLE 3. *STAPHYLOCOCCUS AUREUS* PRESENT IN WOUNDS FOLLOWING TREATMENT WITH MICROCAPSULES A382-140-1, UNLOADED MICROCAPSULES, OR UNENCAPSULATED AMPICILLIN

DAYS POST TREATMENT	MICROCAPSULES	UNLOADED MICROCAPSULES	UNENCAPSULATED AMPICILLIN
(AVERAGE NUMBER* OF ORGANISMS PER GRAM OF TISSUE <sup>†</sup> )			
2	$6.3 \pm 2.6 \times 10^6$	$1.3 \pm 1.5 \times 10^7$	$4.2 \pm 2.4 \times 10^6$
6	$4.7 \pm 2.2 \times 10^5$	$1.0 \pm 5.5 \times 10^6$	$5.3 \pm 7.0 \times 10^6$
8	$7.3 \pm 3.3 \times 10^4$	$4.5 \pm 9.7 \times 10^8$	$5.4 \pm 7.5 \times 10^6$
14	$9.4 \pm 0.6 \times 10^{3§}$	$4.7 \pm 9.8 \times 10^{5×}$	$1.8 \pm 3.2 \times 10^6$

\* Mean  $\pm$  Standard Deviation, n=5

†  $6.0 \times 10^9$  *Staphylococcus aureus* inoculated/wound

§ 3 of 5 wounds sterile

× Lowered due to competitive inhibition by superinfecting gram negative rods

When a regression curve was drawn using the data shown (log concentration of bacteria vs time) the linear component approached significance. This component represented the steady decrease in bacteria/gram of tissue observed in rats treated with microcapsules A382-140-1. When one outlier was removed from each group, significance ( $p < 0.05$ ) was observed.

Report last  
 revised  
 by



SIAM Security  
 Classification  
 Date

SETTERSTROM, TICE, LEWIS, and MEYERS


TABLE 4. AVERAGE NUMBER OF *STAPHYLOCOCCUS AUREUS*/GRAM OF TISSUE PRESENT IN WOUNDS TREATED WITH DECREASING AMOUNTS OF MICROENCAPSULATED AMPICILLIN (A681-31-1)

Bacteria Per Gram of Tissue (mean  $\pm$  S.D., \* n=5)

GROUP	MICROCAPSULE DOSE (g)/WOUND	TOTAL AMPICILLIN AVAILABLE (mg)	48 HOURS		7 DAYS		14 DAYS	
			Bacteria Per Gram of Tissue		Bacteria Per Gram of Tissue		Bacteria Per Gram of Tissue	
A	0.5	90.50	3.7 $\pm$ 1.4X10 <sup>6</sup>		7.7 $\pm$ 10.0X10 <sup>3</sup>		0	
B	0.25	45.25	5.9 $\pm$ 10.0X10 <sup>3</sup>		8.2 $\pm$ 14.0X10 <sup>3</sup>		0	
C	0.10	18.10	2.1 $\pm$ 3.0X10 <sup>6</sup>		1.8 $\pm$ 3.4X10 <sup>4</sup>		0	
D	0.05	9.05	6.8 $\pm$ 4.7X10 <sup>5</sup>		4.4 $\pm$ 3.8X10 <sup>4</sup>		0	
E	0.00	0.00	1.8 $\pm$ 1.8X10 <sup>6</sup>		3.5 $\pm$ 4.3X10 <sup>6</sup>		6.4 $\pm$ 9.6X10 <sup>5</sup>	

\*Standard Deviation  
 1.5X10<sup>10</sup> *S. aureus* inoculated/wound

SIAM Security Classification  
 Date



SETTERSTROM, TICE, LEWIS, and MEYERS

Figure 1. Ampicillin Concentrations Detected in Homogenized Muscle Tissue ( Deep ) at 2, 7, and 14 Days Following Wound Site Treatment with Microencapsulated Ampicillin ( A 681-31-1 )





Figure 2. Ampicillin Concentrations Detected in Serum at 2, 4, 7, and 14 Days Following the Application of Microencapsulated Ampicillin ( Batch A 681-31-1 ) to Infected Wounds



OM, TICE, LEWIS, and MEYERS

#### REFERENCES

- ..., J. F.: "Infection" IN: *Fundamentals of Wound Management*.  
Hunt and J. E. Dunphy (Ed.), Appleton-Century-Crofts, New York,  
1979.
- ..., J. F.: *Surgery* 50:161, 1961.
- ..., C., M. T. Edgerton, M. B. Elliott, L. D. Kurtz, and  
Edlich. *Am J Surg* 127:564, 1974.
- ..., L. E. and J. R. Border. *J Trauma* 10:1078, 1970.
- ..., C., J. Vernick, R. L. Simmons, and T. Motsumoto: *Mil*  
134:13, 1969.
- ..., E. A.: *Brit J Surg* 54:324, 1967.
- ..., D. J.: *Arch Surg* 100:589, 1970.
- ..., N.G., R. S. Howell, and M. Babich: *Arch Surg* 97:365, 1968.
- ..., J.W., G. L. Leonard, and J. L. Ochsner. *Surgery* 67:836,
- ..., D. E., B. Perez, J. Beasley III, W. E. Larson, and W. R.  
...: *Oral Surg* 37:142, 1974.
- ..., T. R., W. E. Meyers, D. H. Lewis, D. R. Cowser, and J. A.  
...: 8th International Symposium on Controlled Release of  
... Materials, Abstracts, 1981.
- ..., V.: *J Labeled Comp* 8:687, 1972.
- ..., D.H., W. E. Meyers, T. R. Tice, and T. O. Dappert: Annual  
... Report, 1981. (Unpublished)









